

COHOMOLOGY OF SYMPLECTIC REDUCTIONS OF GENERIC COADJOINT ORBITS

R. F. GOLDIN AND A.-L. MARE

ABSTRACT. Let \mathcal{O}_λ be a generic coadjoint orbit of a compact semisimple Lie group K . Weight varieties are the symplectic reductions of \mathcal{O}_λ by the maximal torus T in K . We use a theorem of Tolman and Weitsman to compute the cohomology ring of these varieties. Our formula relies on a *Schubert basis* of the equivariant cohomology of \mathcal{O}_λ and it makes explicit the dependence on λ and a parameter in $\text{Lie}(T)^* =: \mathfrak{t}^*$.

1. INTRODUCTION

Let K be a compact semisimple Lie group, $T \subset K$ a maximal torus and $\mathfrak{t} \subset \mathfrak{k}$ their Lie algebras. Pick a fundamental chamber in \mathfrak{t}^* and a point λ in the interior. Let \mathcal{O}_λ be the orbit of λ under the coadjoint representation of K on \mathfrak{k}^* . \mathcal{O}_λ is diffeomorphic to the flag variety K/T and it has a naturally occurring symplectic form ω known as the Kirillov-Kostant-Souriau form. The action of T on \mathcal{O}_λ is Hamiltonian, which means that there is an invariant map

$$\Phi : \mathcal{O}_\lambda \rightarrow \mathfrak{t}^*$$

satisfying $\omega(X_\eta, \cdot) = d\Phi^\eta$, where $\eta \in \mathfrak{t}$, X_η the vector field on \mathcal{O}_λ generated by η , and $\Phi^\eta(m) = \Phi(m)(\eta)$ defined by the natural pairing between \mathfrak{t} and \mathfrak{t}^* . We call Φ a *moment map* for this action.

The image of Φ is the convex hull of $W \cdot \lambda$, the Weyl group orbit of λ . Let $\mu \in \Phi(\mathcal{O}_\lambda)$ be a regular value of Φ . We define the *symplectic reduction* at μ by

$$\Phi^{-1}(\mu)/T = \mathcal{O}_\lambda//T(\mu).$$

The goal of this note is to give a presentation of the cohomology¹ ring of $\mathcal{O}_\lambda//T(\mu)$ in terms of the root system of K . We present $H^*(\mathcal{O}_\lambda//T(\mu))$ as a quotient of the T -equivariant cohomology ring $H_T^*(\mathcal{O}_\lambda)$ by a certain ideal. We rely on the following fundamental result.

Date: October 29, 2002.

¹Only cohomology with coefficients in the field \mathbb{Q} of rational numbers will be considered throughout this paper.

Theorem 1.1 (Kirwan). *Let M be a compact symplectic manifold with a Hamiltonian T action, where T is a compact torus. If $\mu \in \mathfrak{t}^*$ is a regular value of Φ , then the restriction map in equivariant cohomology*

$$\kappa : H_T^*(M) \rightarrow H_T^*(\Phi^{-1}(\mu))$$

is surjective.

As the T action is locally free on level sets of the moment map at regular values, $H_T^*(\Phi^{-1}(\mu)) = H^*(M//T(\mu))$. The resulting map $\kappa : H_T^*(M) \rightarrow H^*(M//T(\mu))$ is called the *Kirwan map*. Kirwan's result is of particular importance because the equivariant cohomology can be described in terms of the equivariant cohomology of the fixed point sets of the T action. In the case of isolated fixed points, this is just a sum of polynomial rings.

Theorem 1.2 (Kirwan). *Let M be a compact Hamiltonian T -space. Let M^T denote the fixed point set of the T action. The restriction map*

$$i^* : H_T^*(M) \rightarrow H_T^*(M^T)$$

is injective. In the case that M^T is a finite set of points, $H_T^(M^T) = \bigoplus_{p \in M^T} \mathbb{Q}[x_1, \dots, x_n]$ where $n = \dim T$.*

A presentation of the cohomology ring of the reduced space $M//T(\mu)$ can be obtained by using the following description of the kernel of the Kirwan map, which is due to Tolman and Weitsman [TW]. If α is in $H_T^*(M)$ we denote

$$\text{supp}(\alpha) = \{p \in M^T : \alpha|_p \neq 0\}$$

Fix an arbitrary inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{t}^* .

Theorem 1.3 (Tolman-Weitsman). *The kernel of the Kirwan map κ is the ideal of $H_T^*(M)$ generated by all $\alpha \in H_T^*(M)$ with the property that there exists $\xi \in \mathfrak{t}^*$ such that*

$$\Phi(\text{supp}(\alpha)) \subset \{x \in \mathfrak{t}^* | \langle \xi, x \rangle \leq \langle \xi, \mu \rangle\}.$$

In other words, α is in $\ker \kappa$ if and only if all points of $\text{supp}(\alpha)$ are mapped by Φ to the same side of an affine hyperplane in \mathfrak{t}^ which passes through μ .*

The T -equivariant cohomology ring of the coadjoint orbit $\mathcal{O}_\lambda = K/T$ is well understood. Kostant and Kumar constructed a basis $\{x_w\}_{w \in W}$ of $H_T^*(K/T)$ as a $H_T^*(pt)$ -module, which we refer to as the *Schubert basis* [KK]. Let B be a Borel in $G := K^\mathbb{C}$, and B_- an opposite Borel. For any $v \in W$, let $X_v = \overline{B_-} \tilde{v} B / B$, where \tilde{v} is any choice of lift of $v \in W$ in the normalizer of the torus. These *opposite Schubert varieties* are

T -invariant subvarieties of $G/B \cong K/T$. The basis $\{x_w\}$ is uniquely defined by the property that

$$\int_{X_v} x_w = \delta_{vw}.$$

Theorem 1.2 suggests the importance of knowing how to restrict the classes x_w to fixed points $W \cdot \lambda$. This formula was worked out for general K by S. Billey [Bi]. In particular, it is easy to show that $x_w|_v = 0$ if $v \not\leq w$ in the Bruhat order.² In other words,

$$\text{supp}(x_w) = \{v\lambda : v \leq w\}.$$

To each $\tau \in W$ we can associate the new basis

$$\{x_w^\tau = \tau \cdot x_{\tau^{-1}w}\}_{w \in W},$$

whose elements have the property

$$\text{supp}(x_w^\tau) = \{v\lambda : \tau^{-1}v \leq \tau^{-1}w\}.$$

Let $\lambda_1, \dots, \lambda_l \in \mathfrak{t}^*$ denote the fundamental weights associated to the chosen fundamental chamber of \mathfrak{t}^* . Let $\langle \cdot, \cdot \rangle$ be the restriction to \mathfrak{t}^* of a K -invariant product on \mathfrak{k}^* . Our main result is:

Theorem 1.4. *The cohomology ring $H^*(\mathcal{O}_\lambda/T(\mu))$ is isomorphic to the quotient of $H_T^*(K/T)$ by the ideal generated by*

$$\{x_v^\tau : \text{there exists } j \text{ such that } \langle \lambda_j, \tau^{-1}v\lambda \rangle \leq \langle \lambda_j, \tau^{-1}\mu \rangle\}.$$

Remarks.

1. One can take the description of $H_T^*(K/T)$ (see for instance [Br]) and deduce a precise presentation of the cohomology ring $H^*(\mathcal{O}_\lambda/T(\mu))$ in terms of generators and relations.
2. For $K = SU(n)$ this result was proven by the first author in [Go1].

Acknowledgement. The second author would like to thank Lisa Jeffrey for introducing him to the topic of the paper. Both authors would like to thank her for a careful reading of the manuscript and for suggesting several improvements.

2. PRIMARY DESCRIPTION OF $\ker \kappa$

For any $\xi \in \mathfrak{t}^*$ we denote by f_ξ the corresponding height function on \mathcal{O}_λ ,

$$f_\xi(x) = \langle \xi, x \rangle.$$

Under the pairing between \mathfrak{t}^* and \mathfrak{t} , the function f_ξ is a component of the moment map. In fact, it is well known that f_ξ is Morse-Bott for all

²The class x_w differs from the ξ^w constructed in [KK] by the relationship $x_w := w_0 \cdot \xi^{w_0 w}$, where w_0 is the longest element of W .

$\xi \in \mathfrak{t}^*$. Denote by $C \subset \mathfrak{t}^*$ the fundamental (positive) Weyl chamber, which can be described by

$$C = \{r_1\lambda_1 + \dots + r_l\lambda_l : \text{all } r_j > 0\},$$

and let \overline{C} be its closure.

Lemma 2.1. *Let τ be in W and ξ in $\tau\overline{C}$. If $\tau^{-1}v < \tau^{-1}w$ in the Bruhat order, then $f_\xi(v\lambda) \leq f_\xi(w\lambda)$.*

Proof. The result follows immediately from the fact that if $\xi \in C$, then the unstable manifold of f_ξ through $v\lambda$ with respect to the Kähler metric on

$$\mathcal{O}_\lambda = K/T = G/B$$

is just the Bruhat cell $B \cdot vB/B$ (see for instance [Ko]). \square

The main result of this section is:

Theorem 2.1. *Suppose that $\alpha \in H_T^*(\mathcal{O}_\lambda)$ has the property that*

$$\Phi(\text{supp}(\alpha)) \subset \{x \in \mathfrak{t}^* : \langle \xi, x \rangle \leq \langle \xi, \mu \rangle\}.$$

Then α can be decomposed as

$$\alpha = \sum_{w \in W} a_w^\tau x_w^\tau$$

with $a_w^\tau \in H_T^(pt)$, such that if $a_w^\tau \neq 0$ then*

$$\Phi(\text{supp}(x_w^\tau)) \subset \{x \in \mathfrak{t}^* : \langle \xi, x \rangle \leq \langle \xi, \mu \rangle\}.$$

Proof. Take $\tau \in W$ such that $\xi \in \tau\overline{C}$. Suppose that the decomposition of α with respect to the basis $\{x_w^\tau\}_{w \in W}$ is of the form

$$(1) \quad \alpha = \sum_{w \in W} a_w^\tau x_w^\tau + a_{v_1}^\tau x_{v_1}^\tau + \dots + a_{v_r}^\tau x_{v_r}^\tau,$$

where the first sum contains only w with

$$\langle \xi, w\lambda \rangle \leq \langle \xi, \mu \rangle,$$

whereas

$$\langle \xi, v_j\lambda \rangle > \langle \xi, \mu \rangle, \quad a_{v_j}^\tau \in S(\mathfrak{t}^*), a_{v_j}^\tau \neq 0,$$

for any $1 \leq j \leq l$. We may assume that v_1 has the property that there exists no $j > 1$ with $\tau^{-1}v_1 < \tau^{-1}v_j$. Now let us evaluate both sides of (1) at $v_1\lambda$. Since

$$\langle \xi, w\lambda \rangle \leq \langle \xi, \mu \rangle < \langle \xi, v_1\lambda \rangle,$$

by Lemma 2.1 we must have

$$x_w^\tau|_{v_1\lambda} = 0$$

for any w corresponding to a term in the first sum in (1). It follows that

$$\alpha|_{v_1\lambda} = a_{v_1}^\tau x_{v_1}^\tau|_{v_1\lambda} \neq 0$$

so $v_1\lambda$ is in $\text{supp}(\alpha)$ even though $\langle \xi, v_1\lambda \rangle > \langle \xi, \mu \rangle$. This is a contradiction. \square

3. PROOF OF THE MAIN RESULT

We now prove Theorem 1.4. Let v and τ in W be such that

$$(2) \quad \langle \lambda_j, \tau^{-1}v\lambda \rangle \leq \langle \lambda_j, \tau^{-1}\mu \rangle,$$

for some $1 \leq j \leq l$. We show that x_v^τ is in the kernel of the Kirwan map

$$\kappa : H_T^*(\mathcal{O}_\lambda) \rightarrow H^*(\mathcal{O}_\lambda//T(\mu)).$$

Let $\xi = \tau\lambda_j$ be in $\tau\overline{C}$. Note that if $w \in \text{supp}(x_v^\tau)$, then $\tau^{-1}w \leq \tau^{-1}v$ implies by Lemma 2.1

$$\langle \xi, w\lambda \rangle \leq \langle \xi, v\lambda \rangle \leq \langle \xi, \mu \rangle.$$

Thus $x_v^\tau \in \ker \kappa$.

Now let us consider $\alpha \in H_T^*(K/T)$ with the property that there exists $\xi \in \mathfrak{t}^*$ with

$$\text{supp}(\alpha) \subset \{x \in \mathfrak{t}^* | \langle \xi, x \rangle \leq \langle \xi, \mu \rangle\}.$$

Take $\tau \in W$ such that $\xi \in \tau\overline{C}$. By Theorem 2.2, we can decompose α as

$$(3) \quad \alpha = \sum_{w \in W} a_w^\tau x_w^\tau$$

where a_w^τ can be nonzero only if

$$\text{supp}(x_w^\tau) \subset \{x \in \mathfrak{t}^* | \langle \xi, x \rangle \leq \langle \xi, \mu \rangle\}.$$

In particular, if $a_w^\tau \neq 0$, then

$$(4) \quad \langle \xi, w\lambda \rangle \leq \langle \xi, \mu \rangle.$$

Since ξ is in $\tau\overline{C}$, it can be written as

$$(5) \quad \xi = \tau \sum_{j=1}^l r_j \lambda_j,$$

where all r_j are non-negative. So (4) and (5) imply that there exists $j \in \{1, \dots, l\}$ such that

$$\langle \tau\lambda_j, w\lambda \rangle \leq \langle \tau\lambda_j, \mu \rangle.$$

In other words, each nonzero term in the right hand side of (3) is a multiple of a x_w^τ of the type claimed in Theorem 1.4. \square

Remark. It follows that, in the particular situation of generic coadjoint orbits, in order to cover the whole Tolman-Weitsman kernel of the Kirwan map it is sufficient to consider affine hyperplanes through μ which are perpendicular to vectors of the type $\tau\lambda_j$, with $\tau \in W$ and $j \in \{1, \dots, l\}$. But these are just the hyperplanes parallel to the walls of the moment polytope. This result concerning a “sufficient set of hyperplanes” has been proved by the first author in [Go2], for an *arbitrary* Hamiltonian torus action on a compact manifold.

REFERENCES

- [Bi] S. Billey *Kostant polynomials and the cohomology of G/B* , Duke Math. J. **96** (1999) 205-224.
- [Br] M. Brion *Equivariant cohomology and equivariant intersection theory*, in *Representation Theory and Algebraic Geometry*, Kluwer Acad. Publ. (1998) 1-37.
- [Go1] R. F. Goldin *The cohomology ring of weight varieties and polygon spaces* Adv. in Math. **160** (2001) No. 2 175-204.
- [Go2] R. F. Goldin *An effective algorithm for the cohomology ring of symplectic reductions* Geom. and Func. Anal., Vol. 12 (2002), 567–583.
- [Ki] F. C. Kirwan, *Cohomology of Quotients in Complex and Algebraic Geometry* Mathematical Notes **31**, Princeton University Press, Princeton, N. J. (1984)
- [Ko] R. R. Koehler *Integral homology of real flag manifolds and loop spaces of symmetric spaces* Adv. in Math. **110** (1995) no. 1, 1-46.
- [KK] B. Kostant and S. Kumar, *The nil Hecke ring and cohomology of G/P for a Kac-Moody group G* Adv. in Math. **62** (1986) no. 3, 187-237.
- [TW] S. Tolman and J. Weitsman, *The cohomology rings of abelian symplectic quotients* math.DG/9807173, to appear in Commun. Anal. Geom.

(R. F. Goldin) MATHEMATICAL SCIENCES, GEORGE MASON UNIVERSITY, MS 3F2, 4400 UNIVERSITY DR., FAIRFAX, VA 22030
E-mail address: rgoldin@gmu.edu

(A.-L. Mare) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ONTARIO M5S 3G3, CANADA
E-mail address: amare@math.toronto.edu