

Social Studies 201
Notes for November 21, 2003

Introduction to hypothesis testing – Chapter 9

These notes introduce hypothesis testing, a topic that will occupy the remainder of the semester. These notes describe the steps involved in tests of hypothesis, the method of testing an hypothesis about a population mean, and some more general principles of hypothesis testing.

Introduction. The aim of hypothesis testing is to bring data to bear on some suspicion or claim, or to examine the credibility associated with a specific research result. Using the principles of hypothesis testing, the data can either negate or support the claim, with a particular probability or significance level attached to the result. Hypothesis tests never provide definitive proof or negation of an hypothesis, but the results of a test can provide strong support for the hypothesis, or they can provide evidence that casts doubt on the hypothesis. Each of these results has a probability attached to it and, in that sense, the conclusion of the test is correct only with such a probability, not with certainty.

Hypotheses are quite specific statements about the characteristics of a population. They emerge from more general statements of the following types.

1. **Suspicions.** Hypotheses may emerge from suspicions or doubts that individuals or groups have about some social issue. One current example is related to the environment – there is suspicion, and a fair degree of evidence, that there is global warming. Exactly what is the cause of this is unclear but one suspicion that people on the Prairies may have is that such global warming could result in productive agricultural regions becoming desert-like. Another suspicion often expressed is that some of the foods or liquids we consume are unsafe – in past years there was suspicion that Regina water was unsafe to drink.

Such suspicions are at a fairly general level and may not be associated with a specific hypothesis about a population mean or proportion. One aim of researchers is to translate some of these suspicions into testable hypotheses, so the data researchers produce can be used to either determine whether these suspicions can be supported or whether they are

unfounded.

2. **Claims.** A more specific type of statement could be considered to be a claim. Claims might be distinguished from suspicions in that the individual making the claim has done some investigation of an issue and has some evidence to make such a claim. But evidence may be incomplete or the claim may remain at a fairly general level.

As examples, consider claims about effects of NAFTA, the North America Free Trade Agreement. Some in the business community and government may claim that NAFTA has expanded the number of jobs in Canada and has been of benefit to Canadians. Detractors of NAFTA may claim that it has led to a loss of jobs, reduced sovereignty, and resulted in a worsening of the living standard of Canadians. Each of the participants in such a debate is likely to have some evidence to support their claim.

A researcher interested in testing these claims would find it necessary to establish particular propositions about the effect of NAFTA, adopt some assumptions about how to examine these issues, and translate the claims into specific hypotheses that can be tested using data the researcher has access to or can obtain.

3. **Research results.** A third, and more specific, level of statement is associated with examination, or re-examination, of the research results of others. Very specific research results may have been made by other researchers, and you may wish to test these results by replicating or questioning them. Other researchers may have conducted well constructed surveys or experiments, but you may wish to re-examine the same issue using data from other populations or using a different theoretical perspective or methodology. These lead you to construct specific hypotheses about the characteristics of the populations you are examining. Again, these lead to statements of hypotheses that can be tested using your data, so that your results either support earlier research or cast doubt on these results.

Each of these levels of research involves construction of specific hypotheses about characteristics of populations – characteristics such as the mean or proportion of a population. Researchers need to take these general statements

and translate them into specific statements about population characteristics such as μ or p . The method of examining such specific hypotheses is the subject of these notes.

Summary of the method of hypothesis testing – pp. 559-563

The idea of hypothesis testing is to hypothesize a particular value for a population parameter such as the population mean, standard deviation, or proportion. If the statistic from a sample of the population is relatively close to the hypothesized value of the population parameter, then the hypothesis is not rejected. But if the sample value is distant from the hypothesized value, the hypothesis is rejected and regarded as incorrect. Attached to each of these conclusions is a probability, that is, an hypothesis is not accepted or rejected with certainty but there is always a probability the the conclusion is correct or incorrect.

In order to illustrate how an hypothesis testing is to be conducted, a test of a population mean will be used as a reference. There are several steps that are involved in testing an hypothesis:

1. **State hypotheses.** The first step in hypothesis testing is to clearly define the population parameter such as a population mena and provide a specific statement about parameter. In the case of a population mean μ , the statement is a claim that μ equals some specific value. This is termed the null hypothesis and given the symbol H_0 . The alternative hypothesis is given the symbol H_1 and is a statement that the mean is not equal to the specific value claimed in H_0 . If the specific value is given a symbol M , then the hypotheses to be tested are:

$$\text{Null hypothesis } H_0 : \mu = M$$

$$\text{Alternative hypothesis } H_1 : \mu \neq M$$

The alternative hypothesis is sometimes termed the research hypothesis and may be of a form the $\mu < M$ or $\mu > M$. More about these options later.

In conducting an hypothesis test, a researcher assumes that the null hypothesis H_0 is correct and the whole test is conducted with this assumption. At the conclusion of the test, this null hypothesis is either rejected (when data are inconsistent with this hypothesis) or not rejected (when data are consistent with the hypothesis).

2. **Obtain sample data.** The sample mean \bar{X} , standard deviation s , and sample size n must be available, or produced by the researcher, in order to determine which of the above hypotheses can be supported using the data available.
3. **Distribution of the test statistic.** If an hypothesis about a population mean is examined, the sample mean is a point estimate of the population mean. The sampling distribution of the sample means \bar{X} must be known in order to proceed with the test. If the samples are large random samples, then the central limit theorem can be used to describe the distribution of these sample means. If the sample size is small, but the samples are random samples from a normally distributed population, then the t-distribution describes the distribution of sample means.
4. **Select a level of significance.** Each hypothesis test is associated with a probability that is termed the level of significance. The level of significance is a small probability and is the complement of the confidence level – for example, 95% confidence level corresponds to 0.05 (or 5%) significance. The level of significance denotes an area in the tail or tails of the distribution of the statistic. The confidence level is a large area in the middle of a normal distribution, denoting a large area within which it is hoped that the sample mean lies. The significance level is an area in the tail or extreme portion of the distribution of the statistic. If the sample is in this extreme area, then H_0 is rejected.

As a result, the significance level specifies a probability that the sample value will be far from the hypothesized mean μ . It indicates a probability that the hypothesis could be rejected in error. Further explanation of this is provided later in these notes.

5. **Critical region or region of rejection of H_0 .** Consistent with the hypothesized value for the mean, the distribution of the sample mean,

and the significance level, a region of rejection for the null hypothesis is constructed. This is a region in the tail or tails of the sampling distribution. In the case of a 0.05 significance level, the critical region is all Z -values less than -1.96 or greater than $+1.96$. That is, there is an area of 0.05 in the two tails of the distribution, or 0.025 of the area in each tail, consistent with a Z value of 1.96. It is this disjointed region that is the critical region or region of rejection of H_0 . The area in the middle of the distribution is the area of nonrejection of H_0 .

6. **Conclusion.** Using the 0.05 level of significance, if the sample mean \bar{X} is farther than 1.96 standard deviations from the mean hypothesized in H_0 , then this null hypothesis is rejected. But if the sample mean \bar{X} is not farther than 1.96 standard deviations from the hypothesized mean, then the null hypothesis H_0 is not rejected. In order to test this, the number of standard deviations that the sample mean lies from the hypothesized mean is calculated.

Remember that a Z -value is also a number of standard deviations, and a Z -value can always be obtained with the formula

$$Z = \frac{\text{variable} - \text{mean of variable}}{\text{standard deviation of variable}}.$$

In this case, when the null hypothesis is that $\mu = M$, the Z -value associated with the sample mean \bar{X} is

$$Z = \frac{\bar{X} - M}{\sigma/\sqrt{n}}.$$

That is, the variable is \bar{X} – its mean is hypothesized to be M and the standard deviation of \bar{X} is σ/\sqrt{n} . This Z -value represents the number of standard deviations the sample mean \bar{X} (with standard deviation σ/\sqrt{n}) is from M . If this value of Z is negative and below -1.96 , or positive and above $+1.96$, then it is in the critical region and the null hypothesis can be rejected. But if this Z -value is between -1.96 and $+1.96$, the sample mean is regarded as close enough to the hypothesized population mean μ so that the null hypothesis is not rejected.

Notation and conventions of hypothesis testing

1. **Null and alternative hypothesis.** The specific hypothesis being tested is termed the null hypothesis and given the symbol H_0 . This represents a specific value for a population parameter, such as $\mu = 50$ or, in the case of proportion, $p = 0.4$. A null hypothesis is a statement that the population parameter has a specific value – as a result the null hypothesis ordinarily an equality.

The alternative hypothesis is a statement concerning other possibilities, for example that $\mu \neq 50$. The alternative hypothesis is sometimes called the research hypothesis and is given the symbol H_1 . It may be a two-directional alternative $\mu \neq 50$ or a one-directional alternative such as $\mu < 50$ or $\mu > 50$. Since the alternative hypothesis is that the population characteristic is different than the specific value hypothesized in H_0 , the alternative hypothesis is ordinarily an inequality.

2. **Level of significance.** This is usually given the symbol α , or alpha, the first letter of the Greek alphabet. This is a small value or probability, perhaps 0.10, 0.05, 0.01, or some other small value. When asking questions about hypothesis testing, a significance level may, or may not, be given. If such a level is not given, the 0.05 level is the most commonly adopted level.
3. **Test statistic.** This is a statistic obtained from a sample – statistics such as \bar{X} or \hat{p} . These are often converted into Z -values or t -values.
4. **Critical values and critical region.** This is the region where the null hypothesis is rejected, and is in the extremes of the distribution. In the case of a normal distribution for the sample mean, this is the Z -value associated with the area in the tail or tails of the normal distribution. For example, if $H_0 : \mu = M$ and $H_1 : \mu \neq M$, and if $\alpha = 0.05$, then the critical values are $Z = -1.96$ and $Z = +1.96$. The critical region is a two-tailed region of all Z -values less than -1.96 or greater than +1.96. If the sample mean is more than this number of Z -values from the hypothesized mean, the null hypothesis is rejected. If the sample mean is not in the critical region (i.e. not more than 1.96 standard deviations from the hypothesized mean) then the null hypothesis is not rejected.

Example – mean age of University of Regina undergraduates

Following is a typical question about testing an hypothesis. A specific claim is stated and sample data are used to test the claim. The data cited here come from the *Survey of Student Attitudes and Experiences* conducted in 1998 in Social Studies 306 and available in the file `ssae.sav` at

```
t:\students\public\201\
```

Question. At the 0.05 level of significance, can you conclude that the mean age of University of Regina undergraduates is 23 years of age? A random sample of 698 undergraduates gave a sample mean age of 22.46 years and a standard deviation of 5.60 years. Use these data to test the contention that the mean age is 23 years.

Answer. The above question requests a test of the hypothesis that the population mean is 23. In this case the population mean to be tested is μ , the mean age of all University of Regina undergraduates in 1998. The steps involved in testing this hypothesis are as follows.

1. **Hypotheses.** The claim is that $\mu = 23$. No direction concerning an alternative is stated in the question so the alternative is that $\mu \neq 23$. The null and alternative hypotheses are

$$H_0 : \mu = 23$$

$$H_1 : \mu \neq 23$$

In words, the null hypothesis H_0 is that the mean age of U of R undergraduates is 23 years of age. The alternative hypothesis H_1 is that the mean age is not 23 years of age. That is, the alternative hypothesis is a two-directional inequality, that the null hypothesis is not correct.

At the conclusion of the test, the data will lead the researcher to either supporting the null hypothesis that μ is 23 years or rejecting this hypothesis. The significance level, the sampling distribution of the test statistic, and the value of the test statistic will determine which of these two conclusions results.

2. **Test statistic.** The claim is about μ , the mean age of all U of R undergraduates. The sample mean of the age of undergraduates in the sample, \bar{X} , is the test statistic that will be used here. If \bar{X} is close to $\mu = 23$, the null hypothesis will not be rejected; if \bar{X} is distant from $\mu = 23$, the null hypothesis will be rejected and the alternative hypothesis that $\mu \neq 23$ will be accepted. The question is how far \bar{X} is from the hypothesized mean of 23. This depends on the distribution of \bar{X} and the significance level.
3. **Distribution of the test statistic.** The sample is said to be a random sample of U of R undergraduates, with a sample size of $n = 698$. This is a large random sample so the central limit theorem can be used. As a result,

$$\bar{X} \text{ is Nor} \left(\mu, \frac{\sigma}{\sqrt{n}} \right).$$

That is, the sampling distribution of \bar{X} is normally distributed with mean μ and standard deviation s/\sqrt{n} , where s can be used as an estimate of the population standard deviation σ , since n is large.

4. **Significance level.** The level of significance requested here is 0.05, so this is $\alpha = 0.05$. This is the area in the two tails of a normal distribution, split equally between each tail so that 0.025 of the area is in each tail.
5. **Critical region.** The critical region is the extreme area that is some distance from the hypothesized mean. In this case, the sample means have a normal distribution and $\alpha = 0.05$, with 0.025 of the area in each tail of the distribution. The critical region is thus all $Z < -1.96$ or $Z > +1.96$, so that there is 0.025 of the area in the critical region at each end of the distribution. The region for rejecting H_0 is thus -1.96 standard deviations or more to the left of centre or +1.96 standard deviations or more to the right of centre.

The area in middle of the distribution is closer to the hypothesized mean, so the null hypothesis is not rejected if the sample is in this middle area. This region of nonrejection of H_0 is all Z -values between -1.96 and +1.96.

These two regions can be summarized as follows:

Region of rejection of H_0 is: $Z \leq -1.96$ or $Z \geq +1.96$

Region of nonrejection of H_0 is: $-1.96 \leq Z \leq +1.96$

6. **Conclusion.** In order to determine whether the sample mean \bar{X} is within the critical region or not, it is necessary to determine the distance \bar{X} is from the hypothesized mean μ . This can be determined by obtaining the Z -value associated with the sample mean – that is, how many standard deviations is $\bar{X} = 22.46$ from the hypothesized mean of $\mu = 23$.

The standard deviation of the mean is s/\sqrt{n} so the number of standard deviations, or Z -values, that \bar{X} is from the mean is

$$\begin{aligned} Z &= \frac{\bar{X} - \mu}{s/\sqrt{n}} \\ &= \frac{22.46 - 23}{5.60/\sqrt{698}} \\ &= \frac{-0.54}{5.60/26.420} \\ &= \frac{-0.54}{0.2120} \\ &= -2.548 < -1.96. \end{aligned}$$

That is, the sample mean is 2.548 standard deviations below the hypothesized mean of $\mu = 23$. This is below the critical cut-off point of -1.96, so this Z -value is in the critical region for the test. That is, the sample mean is -2.548 standard deviations from the hypothesized mean of 23, a great distance, and one that is extreme enough to be in the left 0.025 of the distribution.

Since this Z -value is in the critical region, the conclusion of the test is to reject the null hypothesis H_0 and accept the alternative hypothesis H_1 . This conclusion is made at the $\alpha = 0.05$ level of significance.

The conclusion is that the mean age of U of R undergraduates is not 23 years of age. At the 0.05 level of significance, there is evidence that the mean age is not 23 years, but a different age. The sample yielded a mean age of 22.46, less than 23, so from this the evidence is that the mean age of U of R undergraduates is likely below 23 years of age.