

Non-realizable 2-stage Π -algebras

Hans-Joachim Baues¹ Martin Frankland^{*2}

¹Max-Planck-Institut für Mathematik Bonn

²University of Illinois at Urbana-Champaign

AMS Session on Homotopy Theory
2012 Joint Mathematics Meetings
Boston

Outline

1 Background

2 Criterion for realizability

3 Stable case

1

Background

2

Criterion for realizability

3

Stable case

π_* -modules and Π -algebras

- $X =$ spectrum or space
 $\pi_* X =$ graded group with (primary) **homotopy operations**.
- $X =$ spectrum
 $\pi_* X$ is a $(\pi_* S)$ -module, where $S :=$ sphere spectrum.
 $\alpha \in \pi_k S \rightsquigarrow$ Precomposition $\alpha^*: \pi_n X \rightarrow \pi_{n+k} X$

$$S^{n+k} \xrightarrow{\alpha} S^n \xrightarrow{X} X$$

- $X =$ connected space
 $\pi_* X$ is a Π -algebra.
 $\alpha \in \pi_{n+k} S^n \rightsquigarrow$ Precomposition $\alpha^*: \pi_n X \rightarrow \pi_{n+k} X$

$$S^{n+k} \xrightarrow{\alpha} S^n \xrightarrow{X} X$$

Also π_1 -action and Whitehead products $\pi_p \times \pi_q \rightarrow \pi_{p+q-1}$.

Π -algebra \approx graded group with additional structure which looks like the homotopy groups of a space.

Definition

- $\Pi :=$ full subcategory of the homotopy category of pointed spaces consisting of finite wedges of spheres $\vee S^{n_i}$, $n_i \geq 1$.
- **Π -algebra** := product-preserving functor $\pi: \Pi^{\text{op}} \rightarrow \mathbf{Set}$.

Example

$\pi_* X = [-, X]_*$ for a pointed space X .

Notation: Write $\pi_n := \pi(S^n)$.

Realization and classification

Realization Problem

Given a Π -algebra π , is there a space X satisfying $\pi_* X \simeq \pi$ as Π -algebras?

Classification Problem

If π is realizable, can we classify all realizations?

This project only deals with the **realization problem**.

Remark

A Π -algebra in the stable range is realizable if and only if the corresponding π_* -module is realizable.

2-stage case

Simplest Π -algebras: Only one non-trivial group π_n .

Answer: Always realizable (uniquely), by an Eilenberg-Maclane space $K(\pi_n, n)$.

Next simplest case: Only 2 non-trivial groups π_n, π_{n+k} . Assume $n \geq 2$.

Answer: **Not** always realizable...

Goals

- ① Find necessary and sufficient conditions for a 2-stage Π -algebra to be realizable.
- ② Provide non-realizable examples.

Warm-up

Case $k = 1 \rightsquigarrow$ Always realizable (classic).

Case $k = 2 \rightsquigarrow$ Always realizable (a bit of work).

1 Background

2 Criterion for realizability

3 Stable case

Homotopy operation functors

Reference: Baues, Goerss (2000).

Idea: Encode the Π -algebra data inductively.

Notation

$\Pi\mathbf{Alg}_n^k :=$ full subcategory consisting of Π -algebras concentrated in degrees $n, n+1, \dots, n+k$.

Data consists of abelian groups and structure maps

$$\pi_n$$

$$\eta_1 : \Gamma_n^1(\pi_n) \rightarrow \pi_{n+1}$$

$$\eta_2 : \Gamma_n^2(\pi_n, \eta_1) \rightarrow \pi_{n+2}$$

...

$$\eta_k : \Gamma_n^k(\pi_n, \eta_1, \dots, \eta_{k-1}) \rightarrow \pi_{n+k}.$$

Homotopy operation functors

Postnikov truncation $\Pi\mathbf{Alg}_n^k \rightarrow \Pi\mathbf{Alg}_n^{k-1}$ has a left adjoint L .
 Γ_n^k is the composite

$$\Pi\mathbf{Alg}_n^{k-1} \xrightarrow{L} \Pi\mathbf{Alg}_n^k \xrightarrow{(n+k)^{\text{th}} \text{ group}} \mathbf{Ab}.$$

Example

$$\Gamma_n^1(\pi_n) = \begin{cases} \Gamma(\pi_n) & \text{for } n = 2 \\ \pi_n \otimes_{\mathbb{Z}} \mathbb{Z}/2 & \text{for } n \geq 3. \end{cases}$$

and $\eta_1: \Gamma_n^1(\pi_n) \rightarrow \pi_{n+1}$ is precomposition by the Hopf map
 $\eta: S^{n+1} \rightarrow S^n$.

Homotopy operation functors

A 2-stage Π -algebra consists of the data

$$\begin{aligned} \pi_n \\ \eta_k : \widetilde{\Gamma}_n^k(\pi_n) := \Gamma_n^k(\pi_n, 0, \dots, 0) &\rightarrow \pi_{n+k}. \end{aligned}$$

Notation

$Q_{k,n} :=$ indecomposables of $\pi_{n+k}(S^n)$

In the stable range $k \leq n - 2$, we have $Q_{k,n} = Q_k^S$
($Q_*^S :=$ indecomposables of the graded ring π_*^S).

Proposition

Assuming $k \neq n - 1$, we have

$$\widetilde{\Gamma}_n^k(\pi_n) = \pi_n \otimes_{\mathbb{Z}} Q_{k,n}.$$

In particular, in the stable range we have $\widetilde{\Gamma}_n^k(\pi_n) = \pi_n \otimes_{\mathbb{Z}} Q_k^S$.

Whitehead's exact sequence

$$\dots \rightarrow H_{n+1}X \xrightarrow{b} \Gamma_n X \xrightarrow{i} \pi_n X \xrightarrow{h} H_n X \xrightarrow{b} \Gamma_{n-1} X \rightarrow \dots$$

$h = \text{Hurewicz map}$

$$\Gamma_i X = \text{im}(\pi_i X^{(i-1)} \rightarrow \pi_i X^{(i)})$$

Natural transformation γ :

$$\begin{array}{ccc} \Gamma_n^k(\pi_n, \eta_1, \dots, \eta_{k-1}) & & \\ \downarrow \gamma & \searrow \eta_k & \\ \Gamma_{n+k} X & \xrightarrow{i} & \pi_{n+k} X \end{array}$$

Main result: Criterion for realizability

Theorem (Baues,F.)

The 2-stage Π -algebra given by $\eta_k : \widetilde{\Gamma}_n^k(\pi_n) \rightarrow \pi_{n+k}$ is realizable if and only if the map η_k factors through the map $\gamma_{K(\pi_n, n)}$.

$$\begin{array}{ccc} & \Gamma_{n+k} K(\pi_n, n) & \\ \nearrow \gamma_{K(\pi_n, n)} & \nearrow & \downarrow \\ \widetilde{\Gamma}_n^k(\pi_n) & \xrightarrow{\eta_k} & \pi_{n+k} \end{array}$$

Key ingredient: Theorem on the realizability of the Hurewicz map [Baues].

Remark: $\Gamma_{n+k} K(\pi_n, n) \cong H_{n+k+1} K(\pi_n, n)$.

Main result

Corollary

Fix $n \geq 2$ and $k \geq 1$. Then an abelian group π_n has the property that “every Π -algebra concentrated in degrees $n, n+k$ with prescribed group π_n is realizable” if and only if the map

$$\gamma_{K(\pi_n, n)} : \widetilde{\Gamma_n^k}(\pi_n) \rightarrow \Gamma_{n+k} K(\pi_n, n)$$

is split injective.

1 Background

2 Criterion for realizability

3 Stable case

A simplification

In the stable range ($k \leq n - 2$), the map $\gamma_{K(\pi_n, n)}$ becomes

$$\begin{aligned}\gamma_{K(\pi_n, n)}: \pi_n \otimes_{\mathbb{Z}} Q_k^S &\rightarrow H_{n+k+1} K(\pi_n, n) \\ &\cong (H\mathbb{Z})_{k+1}(H\pi_n) \\ &\cong (H\pi_n)_{k+1}(H\mathbb{Z}).\end{aligned}$$

Proposition

$\gamma_{K(\pi_n, n)}$ factors through the summand $\pi_n \otimes_{\mathbb{Z}} H\mathbb{Z}_{k+1} H\mathbb{Z} \hookrightarrow (H\pi_n)_{k+1} H\mathbb{Z}$.

Upshot

Take $\pi_n = \mathbb{Z}$ and study $\gamma_{K(\mathbb{Z}, n)}: Q_k^S \rightarrow H\mathbb{Z}_{k+1} H\mathbb{Z}$.

Non-realizable example

First few stable homotopy groups of spheres π_*^S and their indecomposables Q_*^S .

k	π_k^S	Q_k^S
0	\mathbb{Z}	\mathbb{Z}
1	$\mathbb{Z}/2\langle\eta\rangle$	$\mathbb{Z}/2\langle\eta\rangle$
2	$\mathbb{Z}/2\langle\eta^2\rangle$	0
3	$\mathbb{Z}/24 \simeq \mathbb{Z}/8\langle\nu\rangle \oplus \mathbb{Z}/3\langle\alpha\rangle$	$\mathbb{Z}/12 \simeq \mathbb{Z}/4\langle\nu\rangle \oplus \mathbb{Z}/3\langle\alpha\rangle$
4	0	0
5	0	0
6	$\mathbb{Z}/2\langle\nu^2\rangle$	0

Non-realizable example

Look at stem $k = 3$.

Proposition

Let $n \geq 5$. The (stable) Π -algebra concentrated in degrees $n, n+3$ given by $\pi_n = \mathbb{Z}$ and $\pi_{n+3} = \mathbb{Z}/4$ with structure map

$$\eta_3: \pi_n \otimes_{\mathbb{Z}} Q_3^S \cong \mathbb{Z}/4\langle\nu\rangle \oplus \mathbb{Z}/3\langle\alpha\rangle \rightarrow \mathbb{Z}/4$$

sending ν to 1 is not realizable.

Proof.

$$H\mathbb{Z}_4 H\mathbb{Z} \simeq \mathbb{Z}/6$$

$\gamma: Q_3^S \simeq \mathbb{Z}/4\langle\nu\rangle \oplus \mathbb{Z}/3\langle\alpha\rangle \rightarrow H\mathbb{Z}_4 H\mathbb{Z}$ sends 2ν to 0. □

Infinite families of indecomposables in π_*^S ?

Good candidates: Greek letter elements (suitable representatives in π_*^S).

Proposition

For every prime $p \geq 3$, the first alpha element $\alpha_1^{(p)} \in Q_{2p-3}^S$ is **not** in the kernel of γ .

Infinite families

Question

What about higher alpha elements $\alpha_i^{(p)} \in Q_{2i(p-1)-1}^S$?

Question

What about beta elements $\beta_i^{(p)}$ ($p \geq 5$)?

Conjecture

Most beta elements $\beta_i^{(p)}$ are in the kernel of γ .

This would provide an infinite family of non-realizable 2-stage (stable) Π -algebras.

Thank you!

franklan@illinois.edu