

This assignment is due at the beginning of class on Monday, April 10, 2006. You must submit all problems that are marked with an asterisk (*).

- 1.** Suppose that Y_1, \dots, Y_n are independent and identically distributed with density function

$$f(y|\theta) = \theta \exp(-\theta y)$$

where $y > 0$ and $\theta > 0$. As usual, let \bar{Y} denote the sample mean given by

$$\bar{Y} = \frac{1}{n} \sum_{i=1}^n Y_i.$$

It is known that the maximum likelihood estimator of θ is $\hat{\theta}_{\text{MLE}} = 1 / \bar{Y}$.

- (a) Consider testing $H_0 : \theta = \theta_0$ against $H_A : \theta \neq \theta_0$. Verify that the rejection region for the generalized likelihood ratio test of these hypotheses is of the form

$$\{\bar{Y} \exp(-\theta_0 \bar{Y}) \leq C\}$$

for some suitable constant C .

- (b) Suppose, to be specific, that $\theta_0 = 1$, and that a random sample of size $n = 10$ is conducted. If the observed data yield $\bar{Y} = 1.25$, perform the generalized likelihood ratio test at the approximate significance level $\alpha = 0.10$.

- 2.** Suppose that Y_1, \dots, Y_n are independent and identically distributed with density function

$$f(y|\theta) = \theta^2 y \exp(-\theta y)$$

where $y > 0$ and $\theta > 0$. As usual, let \bar{Y} denote the sample mean given by

$$\bar{Y} = \frac{1}{n} \sum_{i=1}^n Y_i.$$

It is known that the maximum likelihood estimator of θ is $\hat{\theta}_{\text{MLE}} = 2 / \bar{Y}$.

- (a) Consider testing $H_0 : \theta = 1$ against $H_A : \theta \neq 1$. Verify that the generalized likelihood ratio for this hypothesis testing problem is

$$\Lambda = \left(\frac{\bar{Y}}{2}\right)^{2n} \exp(2n - n\bar{Y}).$$

- (b) Suppose that a random sample of size $n = 5$ is conducted and the observed data yield $\bar{Y} = 1.0$. Perform the generalized likelihood ratio test at the approximate significance level $\alpha = 0.05$.

- 3.** Do the following exercises from Wackerly, et al.

- #10.79 (a), (b), page 515
- #10.83 (a), page 515