

(7.11) To see that  $A$  is a Borel set write  $A$  as

$$A = \{x_0\} = \{(-\infty, x_0) \cup (x_0, \infty)\}^c. \quad (*)$$

Since open intervals are Borel, so too are unions of open intervals, as are complements of unions of open intervals. Using (\*) and elementary properties of the Riemann integral, we have

$$\begin{aligned} P(\{x_0\}) &= \int_{(-\infty, \infty)} \mathbb{1}_{\{x_0\}}(x) f(x) dx \\ &= \int_{(-\infty, x_0)} \mathbb{1}_{\{x_0\}}(x) f(x) dx + \int_{\{x_0\}} \mathbb{1}_{\{x_0\}}(x) f(x) dx + \int_{(x_0, \infty)} \mathbb{1}_{\{x_0\}}(x) f(x) dx \\ &= \int_{(-\infty, x_0)} 0 \cdot f(x) dx + \int_{\{x_0\}} 1 \cdot f(x) dx + \int_{(x_0, \infty)} 0 \cdot f(x) dx \\ &= 0 + \int_{x_0}^{x_0} 1 \cdot f(x) dx + 0 \\ &= 0 \end{aligned}$$

so that  $A$  is a null set for  $P$ .

(7.12) Suppose that  $B$  is countable. Enumerate the elements of  $B$  as  $B = \{x_1, x_2, \dots\}$ . Thus writing  $B = \bigcup_{i=1}^{\infty} \{x_i\}$  expresses  $B$  as a disjoint union. Since  $P$  is a probability, we know that

$$P(B) = P\left(\bigcup_{i=1}^{\infty} \{x_i\}\right) = \sum_{i=1}^{\infty} P(\{x_i\}).$$

But as proved in Exercise 7.11,  $P(\{x_i\}) = 0$  for each  $i$  so that  $P(B) = 0$  as well.

(7.13) If  $P$  and  $B$  are as in Exercise 7.12, and  $P(A) = 1/2$ , then since  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$  we conclude

$$\begin{aligned} P(A \cup B) &= \int_{-\infty}^{\infty} \mathbb{1}_{A \cup B}(x) f(x) dx = \int_{-\infty}^{\infty} \mathbb{1}_A(x) f(x) dx + \int_{-\infty}^{\infty} \mathbb{1}_B(x) f(x) dx - \int_{-\infty}^{\infty} \mathbb{1}_{A \cap B}(x) f(x) dx \\ &= 1/2 + 0 - \int_{-\infty}^{\infty} \mathbb{1}_{A \cap B}(x) f(x) dx. \end{aligned}$$

We know from Exercise 7.12 that  $B$  is a null set for  $P$  (and that  $B$  is actually a Borel set). Since we are told that  $A$  is an event, we can conclude that  $A \cap B$  is an event as well. Since  $A \cap B \subseteq B$ , we see that  $P(A \cap B) \leq P(B) = 0$  so that

$$\int_{-\infty}^{\infty} \mathbb{1}_{A \cap B}(x) f(x) dx = 0,$$

and therefore  $P(A \cup B) = 1/2$ , as required.

An alternative solution is as follows. Since  $B$  as given by Exercise 7.12 is a Borel set, and since  $A$  is assumed to be an event, we know that  $A \cup B$  is also an event. It now follows that  $P(A \cup B) = 1/2$  since  $A \subseteq B$  implies

$$\frac{1}{2} = P(A) \leq P(A \cup B) \leq P(A) + P(B) = \frac{1}{2} + 0 = \frac{1}{2}.$$

(7.14) Suppose that  $A_1, A_2, \dots$  is a sequence of null sets. This means that there exist sets  $C_1, C_2, \dots$  with  $A_i \subseteq C_i$  and  $P(C_i) = 0$  for each  $i$ . Let

$$C = \bigcup_{i=1}^{\infty} C_i$$

so that

$$B = \bigcup_{i=1}^{\infty} A_i \subseteq C.$$

Since

$$P(C) = P\left(\bigcup_{i=1}^{\infty} C_i\right) \leq \sum_{i=1}^{\infty} P(C_i) = 0$$

we conclude that  $B$  is a null set for  $P$ .

(7.15) Suppose that  $E(|X|) = 0$ . To show that  $X = 0$  except possibly on a null set means to show that  $P(X = 0) = 1$ . We will prove  $P(X = 0) = 1$  by deriving a contradiction. Suppose, to the contrary, that  $P(X = 0) < 1$ . Then, there exists some  $a > 0$  such that  $P(|X| \geq a) > 0$ . However, by Markov's inequality (Corollary 5.1), we have that for every  $a > 0$ ,

$$P(|X| \geq a) \leq \frac{E(|X|)}{a} = 0$$

since  $E(|X|) = 0$  by assumption. Hence, for every  $a > 0$ , we have  $P(|X| \geq a) = 0$ , and we conclude  $P(|X| > 0) = 0$ , or in other words,  $P(X = 0) = 1$ .

It is not possible to conclude in general that  $X = 0$  everywhere. As a simple example, suppose that  $\Omega = \{0, 1\}$  and let  $P$  be the Dirac mass at the point 0. (See Example 2 on page 42.) It then follows that the random variable  $X : \Omega \rightarrow \{0, 1\}$  whose law (or distribution) is  $P$  has  $P(X = 0) = 1$  and  $P(X = 1) = 0$  so that  $E(|X|) = 0$ , even though  $X \neq 0$  everywhere (i.e.,  $X(\omega) \neq 0$  for some  $\omega \in \Omega$ ).

(7.17) A direct application of Corollary 7.1 gives

$$(a) \quad P\left(\left(-\frac{1}{2}, \frac{1}{2}\right)\right) = F\left(\frac{1}{2}-\right) - F\left(-\frac{1}{2}\right) = 1/4 - 0 = 1/4,$$

$$(b) \quad P\left(\left(-\frac{1}{2}, \frac{3}{2}\right)\right) = F\left(\frac{3}{2}-\right) - F\left(-\frac{1}{2}\right) = (1/4 + 1/2) - 0 = 3/4,$$

$$(c) \quad P\left(\left(\frac{2}{3}, \frac{5}{2}\right)\right) = F\left(\frac{5}{2}-\right) - F\left(\frac{2}{3}\right) = (1/4 + 1/2 + 1/4) - (1/4) = 3/4,$$

$$(d) \quad P([0, 2]) = F(2-) - F(0-) = (1/4 + 1/2) - 0 = 3/4,$$

$$(e) \quad P((3, \infty)) = 1 - P((-\infty, 3]) = 1 - F(3) = 1 - (1/4 + 1/2 + 1/4) = 0.$$

(7.18) In order to prove that the function  $F$  given by

$$F(x) = \sum_{i=1}^{\infty} 2^{-i} \mathbb{1}_{[\frac{1}{i}, \infty)}(x)$$

is a distribution function of a probability on  $\mathbb{R}$  we use Theorem 7.2. Clearly,  $F(x) = 0$  for all  $x \leq 0$ , so that

$$\lim_{x \rightarrow -\infty} F(x) = 0.$$

Moreover, for all  $x \geq 1$ ,

$$F(x) = \sum_{i=1}^{\infty} 2^{-i} \mathbb{1}_{[\frac{1}{i}, \infty)}(x) = \sum_{i=1}^{\infty} 2^{-i} = 1$$

so that

$$\lim_{x \rightarrow \infty} F(x) = 1.$$

Suppose that  $0 < x < y$ . If  $x \geq 1/i$  for some  $i$ , then necessarily  $y > 1/i$  since  $y > x$ . In particular,  $\mathbb{1}_{[\frac{1}{i}, \infty)}(x) \leq \mathbb{1}_{[\frac{1}{i}, \infty)}(y)$  for all  $-\infty < x < y < \infty$  so that  $F$  is non-decreasing. We have already shown that  $F(x) = 0$  for all  $x \leq 0$  and that  $F(x) = 1$  for all  $x \geq 1$  so that  $F$  is necessarily right continuous on  $(-\infty, 0) \cup [1, \infty)$ . We must still show that  $F$  is right continuous for all  $x \in [0, 1)$ . Notice that  $F$  is a step function for  $0 \leq x < 1$  with jumps at the points  $x = 1/i$ ,  $i = 2, 3, \dots$ . It is therefore clear that  $F$  is continuous on each open interval  $((i+1)^{-1}, i^{-1})$ , for  $i = 1, 2, \dots$ . Suppose that  $x = 1/i$  for some  $i = 1, 2, \dots$ . Then, for all  $y$  with  $1/(i-1) > y > 1/i$  we have  $F(y) = F(1/i)$  so that

$$F(x+) = F(1/i+) = \lim_{y \rightarrow 1/i+} F(y) = \lim_{y \rightarrow 1/i+, y > 1/(i-1)} F(1/i) = F(1/i).$$

It remains to show that  $F$  is right continuous at 0; that is, we must show

$$F(0+) = \lim_{y \rightarrow 0+} F(y) = 0. \quad (\dagger)$$

To prove  $(\dagger)$ , we show that for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that  $F(y) < \varepsilon$  whenever  $y < \delta$ . Let  $\varepsilon > 0$  be arbitrary. Then there exists an  $i_0 \in \mathbb{N}$  such that  $\varepsilon \geq 2^{-i_0}$ . Let  $\delta = 1/i_0$  so that  $y < 1/i_0$ . Thus, by the right-continuity of  $F$ ,

$$F(y) \leq F(1/i_0) = \sum_{i=1}^{\infty} 2^{-i} \mathbb{1}_{[\frac{1}{i}, \infty)}(1/i_0) = \sum_{i=i_0}^{\infty} 2^{-i} = 1 - \sum_{i=1}^{i_0-1} 2^{-i} = 1 - \frac{1 - 2^{-i_0}}{1 - 1/2} = 2^{-i_0} \leq \varepsilon.$$

Thus, by Theorem 7.2,  $F$  is the distribution function of a probability on  $\mathbb{R}$ .

Finally, a direct application of Corollary 7.1 gives

$$(a) \ P([1, \infty)) = 1 - F(1-) = 1 - \sum_{i=2}^{\infty} 2^{-i} \mathbb{1}_{[\frac{1}{i}, \infty)} = 1 - \sum_{i=2}^{\infty} 2^{-i} = 1 - 1/2 = 1/2,$$

$$(b) \ P\left(\left[\frac{1}{10}, \infty\right)\right) = 1 - F\left(\frac{1}{10}-\right) = 1 - \sum_{i=11}^{\infty} 2^{-i} \mathbb{1}_{[\frac{1}{i}, \infty)} = \sum_{i=1}^{10} 2^{-i} = \frac{1 - 2^{-11}}{1 - 1/2} - \frac{1}{2} = 1 - 2^{-10},$$

$$(c) \ P(\{0\}) = F(0) - F(0-) = 0 - 0 = 0,$$

$$(d) P\left(\left[0, \frac{1}{2}\right)\right) = F\left(\frac{1}{2}-\right) - F(0-) = \sum_{i=3}^{\infty} 2^{-i} \mathbb{1}_{\left[\frac{1}{i}, \infty\right)} - 0 = 1 - \sum_{i=1}^2 2^{-i} = 1 - (1/2 + 1/4) = 1/4,$$

$$(e) P((-\infty, 0)) = F(0-) - 0 = 0,$$

$$(f) P((0, \infty)) = 1 - P((-\infty, 0]) = 1 - F(0) = 1 - 0 = 1.$$